
Association Between Acute Kidney Injury, Delirium, & Outcomes in Patients with Critical Illness: A Systematic Review & Meta-Analysis





Prit Kusirisin, MD, PhD^{1,2},








University of Alberta & Alberta Health Services, Edmonton, Alberta, Canada
 Division of Nephrology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand



Introduction: AKI & Delirium in Critically III Patients



Direct effect

- Cytokines
- DAMPs, PAMPs
- Oxidative stress

Indirect effect

- Immune cells
- Sympathetic or parasympathetic activation

Possible effect

CNS, central nervous system;

DAMPs, damage-associated molecular patterns; PAMPs, pathogen-associated molecular patterns

Proposed Mechanisms of AKI-Associated Delirium & Potential Cellular Drivers of Delirium Impaired clearance Inflammation ↑ vascular Electrolyte < of toxins & drugs permeability imbalances including sedatives **Acute kidney injury** ↓ central Periphery CNS Disruption dopamine turnover Fluid accumulation Blood of BBB Cytokine endotoxin/ cytokines release **BBB** Cerebral edema activation Microglial TNF-α IL-6 activation IL-1α Hippocampal IL-1β injury Brain Neuronal cytokines dysfunction Astrocyte

activation

Knowledge gap (review question)

& injury

No systematic evaluation describing clinical association between AKI & delirium among critically ill patients

Objective

BBB

To perform systematic review describing association between AKI & delirium among patients with critical illness

Blood

vesse

Methods

- Study registration: PROSPERO on March 25, 2025 (CRD420251001864)
- Searching
 - Databases: Ovid MEDLINE, Ovid Embase, CINAHL, Scopus, Web of Science Core Collection, Cochrane library, Google scholar
 - Keywords: acute kidney injury, renal replacement therapy (RRT), critical care, critical illness, intensive care, intensive care units, delirium
- 2 reviewers independently screened, selected, extracted studies, & assessed risk of bias using web-based tool Covidence

Inclusion criteria

- Clinical trial or observational cohort
- Study reports occurrences of both AKI & delirium, using validated definitions
- Study focused on population admitted to critical care or intensive care unit (ICU)
- Published between January 2000 to January 2025
- Published in English language

Exclusion criteria


- Case reports or case series
- Pre-clinical or experimental studies

Methods

Population	Critically ill patients with AKI					
Comparison	Develop delirium vs Do not develop delirium					
Outcomes	 Primary: proportion of de novo delirium Secondary: Odds of delirium stratified by AKI severity In-hospital mortality Use of organ support: RRT, mechanical ventilation (MV), vasopressor Length of ICU & hospital stay 					


- Qualitative synthesis: summary characteristics of included studies
- Quantitative synthesis
 - Dichotomous outcomes: pooled proportion, risk ratio (RR), or odds ratio (OR), with 95% CI
 - Continuous outcomes: weighted mean differences (WMD), with 95% CI
 - Pooled meta-analysis: DerSimonian-Laird random-effects model
 - Statistical heterogeneity: visual inspection, Q test <0.10, I2 statistic >50%
 - Publication bias: funnel plot (if >10 studies per outcomes)
 - Statistical significance: p < 0.05
 - Strength of evidence: GRADE tool

STATA
statistical software,
version 17




Results: PRISMA Diagram

- Search complete on March 25, 2025
- Included 18 observational studies
- Study quality
 - Using Newcastle-Ottawa
 Scale
 - Moderate quality (mean 6.7±1.3, range of 4-8)




Characteristics of Included Studies

- 18 studies ≈ 158,694 patients
- Population setting
 - Surgical ICU (10), medical ICU (1), cardiac ICU (1), mixed ICU (5), pediatric ICU (1)
- Definition
 - AKI
 - KDIGO (8), AKIN (1), N/A (9)
 - Delirium
 - CAM-ICU (12), DSM (2),
 CAPD (1), NuDesc (1), N/A (1)




Primary Outcome

Proportion of Delirium




Proportion of delirium among patients with AKI 32% (95% CI, 18-47)




Secondary Outcomes


				Mor	ality		
	Delirium		No delirium		Risk r	atio	Weight
Study	Events	Total	Events	Total	with 95	% CI	(%)
Jackel et al, 2021	50	230	12	153	2.46 [1.35,	4.47]	11.55
Kalyoncuoglu et al, 2020	13	17	13	61	2.47 [1.30,	4.68]	11.10
Koth et al, 2022	28	629	2	279	5.99 [1.44,	24.96]	5.00
Lescot et al, 2013	9	28	30	253	2.29 [1.18,	4.45]	10.89
Lin et al, 2023	19	220	20	472	1.96 [1.06,	3.59]	11.45
Mungan et al, 2020	1	10	0	33 -	8.50 [0.37,	194.86]	1.38
Pauley et al, 2015	39	120	24	470	5.05 [3.14,	8.13]	12.89
Shin et al, 2018	14	40	3	59	5.36 [1.63,	17.65]	6.32
Trongtrakul et al, 2016	159	55	286	390	1.76 [1.56,	1.98]	15.81
Wan et al, 2020	32	142	44	142	0.78 [0.52,	1.17]	13.62
Overall					2.36 [1.61,	3.47]	
Heteroger							
Test of θ _i :	In-hospital mortality						
Test of θ =			_				
	RR :	2.3	6 (9	5%	CI, 1.61-3.47;		
andom-ef		_					
p <0	.01,	- =	79	%; I	noderate certainty		


		F	Receipt o	of Mecha	nical Ventilation			
	Deli	rium	No delirium		Risk ratio Wei	ght		
Study	Events	Total	Events	Total	with 95% CI (%	s)		
Chen et al, 2024	676	1,133	855	3,685	1.98 [1.82, 2.16] 10.	50		
Jackel et al, 2021	193	230	62	153	1.58 [1.25, 2.00] 10.	11		
Koth et al, 2022	320	629	20	279	5.04 [3.27, 7.77] 9.	15		
Lin et al, 2023	220	220	472	472	1.00 [0.89, 1.12] 10.	45		
Marquetand et al, 2021	208	1,285	24	1,791	10.54 [6.94, 15.99] 9.	24		
Pauley et al, 2015	77	120	86	470	2.53 [1.95, 3.28] 10.	00		
Singh et al, 2020	7,550	14,645	9,950	14,645	0.84 [0.82, 0.86] 10.	56		
Tian et al, 2020	93	93	527	527	1.00 [0.86, 1.17] 10.	36		
Wan et al, 2020	113	142	96	142	1.10 [0.89, 1.35] 10.	20		
Yang et al, 2023	27	1,020	510	114,127	5.80 [3.96, 8.50] 9.	43		
Overall					2.08 [1.46, 2.98]			
Heterogeneity								
Test of $\theta_i = \theta_j$	Receipt of MV							
Test of $\theta = 0$:		_						
	RR 2.08 (95% CI, 1.46-2.98;							
p <0.01, $I^2 = 99\%$; low certainty)								
handom-enect	p <(J.U1	, =	99%	; low certainty)			





Secondary Outcomes: Length of Stay





Length of ICU stay
WMD 3.54 days

 $(95\% CI, 1.20-5.87; p < 0.01, I^2 = 99\%; moderate certainty)$

Length of hospital stay

WMD 4.78 days

 $(95\% CI, 3.48-6.09; p < 0.01, I^2 = 91\%; moderate certainty)$

Discussion

Key findings

- Common incidence of delirium (32%) among critically ill patients with AKI
- Stepwise ↑ risk of developing delirium with worsening AKI stage
- Delirium associated with ↑ risk of mortality,
 ↑ receipt of RRT, MV, & vasoactive therapy,
 as well as longer ICU & hospital stays
- Offer new insights by comprehensively synthesizing current evidence on the association between AKI & delirium in critically ill patients

Limitation

- Quality & heterogeneity of included studies
 - Diverse populations
 - Varying definitions of AKI & delirium
 - Regional disparities in healthcare systems & AKI management
- Absence of clear time points for delirium onset relative to AKI, which limits interpretation of causality

Clinical application


- † awareness of incident delirium
- Early detection & management of AKI to mitigate the risk of kidney-brain crosstalk

Conclusion

- Delirium is common among critically ill patients with AKI & is associated with worse clinical outcomes & greater health resource use.
- Strategies to improve the recognition & monitoring of delirium may enhance ICU care & mitigate adverse patient outcomes.

Acknowledgements

Prof. Sean M Bagshaw
Conceive protocol





Assoc. Prof.
M Elizabeth Wilcox



Assoc. Prof.
Oleksa G Rewa



Janice Y Kung
Search strategy



Angela Corsaro
Second reviewer



Thank You for Your Attention NEPHRO MED CMU



